

**Subject:** Scoliosis Surgery  
**Document #:** SURG.00097  
**Status:** Revised

**Publish Date:** 07/06/2022  
**Last Review Date:** 05/12/2022

## Description/Scope

This document addresses surgical treatments for scoliosis, specifically, use of a minimally invasive deformity correction system (ApiFix, Ltd, Misgav Business Park, Israel), vertebral body tethering, and vertebral body stapling. This document does not address spinal fusion for scoliosis treatment.

## Position Statement

### Medically Necessary:

Vertebral body tethering for the treatment of scoliosis is considered **medically necessary** for **all** of the following criteria:

- Curve progression following conservative management (for example, observation, exercise therapy, or bracing); **and**
- Cobb angle 40 to 60 degrees; **and**
- Curve flexibility greater than 30%; **and**
- Skeletal immaturity, defined as either:
  - Risser grade 0 or 1; **or**
  - Sanders Maturity Scale less than or equal to 4.

### Investigational and Not Medically Necessary:

Vertebral body tethering for the treatment of scoliosis is considered **investigational and not medically necessary** when the above criteria are not met and for all other indications.

Use of a minimally invasive deformity correction system for the treatment of scoliosis is considered **investigational and not medically necessary**.

Vertebral body stapling for the treatment of scoliosis is considered **investigational and not medically necessary**.

## Rationale

### Vertebral Body Tethering

Primary treatments for scoliosis include exterior bracing or surgical spinal fusion. Vertebral body tethering has also been proposed as an alternative to bracing in the treatment of scoliosis.

The United States Food and Drug Administration (FDA) has granted Humanitarian Device Exemption (HDE) for vertebral body tethering. The exemption was based on a clinical trial of 56 participants who had spinal tethering

Federal and State law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. The member's contract benefits in effect on the date that services are rendered must be used. Medical Policy, which addresses medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

**Scoliosis Surgery**

---

around 12 years of age. In this clinical trial the participants, on average, had a Cobb angle curve reduction by more than 50%, from 40.4 degrees to 17.6 degrees, at or beyond 24 months post-procedure. Of the 43 participants with a pre-operative Cobb angle of less than 45 degrees, 35 (81.4%) achieved a Cobb angle less than 30 degrees; of the 12 participants with a pre-operative Cobb angle of greater than or equal to 45 degrees, a Cobb angle less than 30 degrees was achieved in 8 (61.5%). The most common complications included back pain, overcorrection of the curve, nausea/vomiting, arm and leg pain, temporary numbness in the chest and hip, and the need for additional surgery. Eight of the participants required an additional surgery to fix overcorrections, cord breakage, development of a new curve in another area of the spine, and slippage in the spine unrelated to the tethering.

A 2014 retrospective review by Samdani and colleagues reported on the 2-year results of 11 participants with scoliosis who underwent vertebral body surgery (8 participants had tethering and 3 participants had tethering and stapling). All of the participants were skeletally immature prior to undergoing surgery. The tethering was done on an average of 7.8 levels. Preoperative thoracic Cobb angle averaged  $44.2 \pm 9.0^\circ$  and corrected to  $20.3 \pm 11.0^\circ$  on first erect, and 2-year Cobb angle  $13.5 \pm 11.6^\circ$ . The preoperative lumbar curve of  $25.1 \pm 8.7^\circ$  showed correction (first erect =  $14.9 \pm 4.9^\circ$  and  $7.2 \pm 5.1^\circ$  at 2 years). Thoracic axial rotation as measured by a scoliometer went from  $12.4 \pm 3.3^\circ$  preoperatively to  $6.9 \pm 3.4^\circ$  at the most recent measurement. There were no neurological, infectious or hardware-related complications noted. Two of the participants required an additional surgery to loosen the tether secondary to overcorrection. One participant had persistent atelectasis.

Samdani and colleagues (2015) reported on retrospective chart review of another cohort of individuals with scoliosis who underwent vertebral body tethering. Following 32 skeletally immature participants with a 1-year follow-up, the participants had tethering of an average of 7.7 levels. The mean preoperative thoracic curve was  $42.8^\circ \pm 8.0^\circ$  which corrected to  $21.0^\circ \pm 8.5^\circ$  on first erect and  $17.9^\circ \pm 11.4^\circ$  at most recent. The pre-operative lumbar curve of  $25.2^\circ \pm 7.3^\circ$  showed correction at first erect  $18.0^\circ \pm 7.1^\circ$  and 1-year correction of  $12.6^\circ \pm 9.4^\circ$ . Preoperative thoracic axial rotation measured  $13.4^\circ$  with correction to  $7.4^\circ$  at the most recent measurement. There were no neurological, infectious or hardware-related complications noted. Overcorrection occurred in three participants who the authors note may require an adjustment of the tether. The authors state “These early results appear promising, but the true benefit of the technique must stand the test of time.”

In a single-center phase 2A pilot study by Wong and colleagues (2019), the authors reported on the use of an anterior ultra-high molecular weight polyethylene tether in 5 children with thoracic scoliosis. Participants were followed for a minimum of 4 years. The preoperative mean thoracic Cobb angle was  $40.1^\circ$ . The degree of correction at 4 years ranged from 0-133.3%. There were 20 adverse events postoperatively, 4 of which were considered to be of moderate severity including pneumonia, distal decompensation, and curve progression. Overcorrection occurred in 3 of the participants, of which 2 required fusion surgery.

A 2020 retrospective review by Hoernschemeyer and colleagues reported on 31 participants with scoliosis who had vertebral body tethering. Two participants were lost to follow-up. The average follow-up was 3.2 years. In this study, outcomes were considered successful where there was a residual curve of less than  $30^\circ$  when the participants were skeletally mature and did not require fusion surgery. A total of 20/27 participants showed a curve magnitude of less than  $30^\circ$ . There were 14 participants found to have broken tethers (5 occurred during the first 2 years, 8 occurred between year 2 and 3, and 4 occurred after the third year of follow-up). Of the 14 participants found to have a broken tether, 7 participants were considered clinically successful, 5 were unsuccessful, and 2 had fusion surgery for continued curve progression. The study is limited by the retrospective design and lack of outcomes

---

Federal and State law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. The member's contract benefits in effect on the date that services are rendered must be used. Medical Policy, which addresses medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

reported by the participants. Two of the participants in this study have not reached skeletal maturity so their clinical success remains unknown.

A 2020 retrospective study by Newton and colleagues compared the outcomes of participants with scoliosis who received vertebral body tethering (n=23) to a matched cohort who were treated with posterior spinal fusion (n=26). The mean follow-up was 3.4 years in the vertebral body tethering group and 3.6 years in the spinal fusion group. Preoperative mean thoracic curve was 53° in the tethering group and 54° in the fusion group. At the final follow-up, the mean thoracic curve was 33° in the tethering group and 16° in the fusion group. In the tethering group, there were 9 revisions and no revisions in the fusion group. Revision procedures occurred at a mean time of 2.3 years postoperatively. Broken tethers were experienced by 12 participants and 3 of the participants had revision due to curve progression from the tether breakage. In the tethering group, 12 participants were considered to have clinical success as evidenced by thoracic curve less than 35° without a secondary spinal fusion. All of the participants in the spinal fusion group had curves of less than 35°.

A multicenter, retrospective review by Miyajima and colleagues (2020) reported on the clinical efficacy of vertebral body tethering in skeletally immature individuals with idiopathic scoliosis. In this study, 57 individuals had vertebral body tethering surgery. The mean follow-up was 40.4 months. The mean preoperative major curve was 51° which improved to a mean of 24.6° following surgery. At one year post-op, the mean major curve was 16.3° and at final follow-up it was 23°. There were 16 complications with 8 individuals who required an additional 9 unplanned revision surgeries. An additional five participants had fusion for insufficient correction and progression of deformity. Most of the outcomes were considered successful, but the outcomes varied due to the 28.1% complication rate and 15.8% of participants requiring further surgery with a mean follow-up of 40.4 months. The authors concluded “a minimum two-year follow-up is not an adequate benchmark for these patients and clearly longer follow-up is required to make any definitive statements about the true value of this technique.”

Baker and colleagues completed a retrospective chart review in 2021 and reported on 17 participants with adolescent idiopathic scoliosis. The objective was 2-4 year outcomes following vertebral body tethering surgery. There were nine curves in 9/17 participants which were considered successful. Preoperative kyphosis averaged 26° in the successful group and 14° in the unsuccessful group. The authors looked at both lumbar and thoracic levels and noted correction was greater for the lumbar region compared to thoracic tethering. There were nine broken tethers and four participants required revision procedures. The authors concluded the technology has potential for those with adolescent idiopathic scoliosis looking for an alternative to spinal fusion surgery, however predicting those who will have successful outcomes is a challenge and long-term outcomes remain unknown.

In 2021 Samdani and colleagues published study results from the FDA Investigational Device Exemption for vertebral body tethering. This industry sponsored, single center study included 57 skeletally immature subjects who underwent vertebral body tethering with 56 participants included in the final analysis. Follow-up lasted for an average of 55.2 months. Mean main thoracic Cobb angle was 40.4° with correction to 19.3° at the first erect measurement and 13.8° at the 2-year follow-up. Most recent follow-up showed mean major thoracic Cobb angle at 18.7°. There were 45/56 subjects with curves less than 30° at the latest follow-up. Mean preoperative proximal thoracic lumbar curve was 25.0° and lumbar curve was 23.7°. At the latest follow-up, mean proximal thoracic curve was 17.9° and lumbar curve was 15.7°. Mean preoperative kyphosis measured 15.5°, 17° postoperatively and 19.6° at the most recent follow-up. There was no significant change in lumbar lordosis from preoperative measurement to the most recent evaluation. Seven of the subjects required revision surgery (5 had tether release for overcorrection

---

Federal and State law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. The member's contract benefits in effect on the date that services are rendered must be used. Medical Policy, which addresses medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

and 2 had tether extension for adding-on). The tether release did not stop the curve overcorrection in one of the subjects and later had posterior spinal fusion surgery.

A 2020 prospective case series by Rushton and colleagues sought to determine the efficacy of vertebral body tethering in 112 skeletally immature subjects with idiopathic scoliosis. Indications for tethering included progressive major main thoracic and/or lumbar curves greater than or equal to 40°. Outcomes were measured preoperatively, radiograph at first erect, 1-year postoperatively, and at most recent follow-up. Clinical success was defined as lack of waiting for or completed surgical fusion and tethered curve less than 35°. Mean follow-up was 37 months. Preoperative mean coronal Cobb curve was 50.8°, corrected to 26.6° at the first erect x-ray. From first erect to 1-year follow-up, mean Cobb curve was 23.1°. There were 36 tethers suspected or confirmed broken postoperatively, 2 of which were replaced by most recent follow-up. The remaining 34 suspected/confirmed broken tethers at follow-up had an increase Cobb angle from 26.1° at 1-year to 35.1° at most recent follow-up. The subjects with intact tethers maintained a mean Cobb angle of 21.7° at 1-year to 21.8° at most recent follow-up. Kyphosis preoperatively was 18.6° which was stable at 1-year then increased to 21.4° at most recent follow-up. At most recent follow-up, 7 subjects had undergone or were waiting to have fusion surgery. The remaining 80 subjects had tethered curve less than 35° were considered clinically successful in this study. There were 25 subjects who had 28 complications including atelectasis, hemothorax, pneumonia, cerebrospinal fluid leaks, additional surgery for bleeding control with 15 subjects requiring 18 revision procedures. Confirmed tether breakage was found in 3 cases. X-rays suggested tether breaks in 33 subjects. The authors state “Further work is needed to examine the variability seen in patient responses to tethering and impact of tether breakage. Prolonged follow up of AVBT patients will be needed until we can understand the true value of this technique.”

A 2021 study by Pehlivanoglu and colleagues reported on the clinical and functional outcomes of comparing vertebral body tethering to posterior spinal fusion. There were 21 subjects who had vertebral body tethering and 22 subjects who had fusion surgery. Average follow-up for the tethering group was 37.1 months and 37.8 months for the fusion group. Functional evaluation performed at the last follow-up appointment was measured by average lumbar range of motion, anterior lateral lumbar bending flexibility, flexor and extensor endurance of trunk, and average motor strength of trunk muscles. Scoliosis Research Society (SRS-22) and Short Form Survey (SF-36) scores were used to evaluate functional outcome and health-related quality of life. Preoperative average major curve magnitude in the tethering group was 48.2° and 48.8 in the fusion group. Average major curve magnitude at the last follow-up was 9.1° in the tethering group and 9.7° in the fusion group. In the tethering group, average flexion lumbar range of motion was 78.2, average extension was 34.6, average lateral bending was 34.4, average rotation was 45.4, average anterior lumbar flexibility was 3.7 and average lateral lumbar flexibility was 22.4. In the fusion group, average flexion lumbar range of motion was 58.1, average extension was 19.4, average lateral bending was 18.3, average rotation was 24.1, average anterior lumbar flexibility was 23.4 and average lateral lumbar flexibility was 11.3. In the tethering group, average flexor trunk endurance was 65.1 and extensor endurance was 60.8 with average motor strength of trunk muscles 4.7. In the fusion group, average flexor trunk endurance was 19.1 and extensor endurance was 28.7 with average motor strength of trunk muscles 3.2. Average pre-operative SRS-22 score in the tethering group was 3.2 and 4.9 at the latest follow-up. Fusion group had average pre-operative SRS-22 score of 3.2 and 3.8 at latest follow-up. Average pre-operative SF-36 scores for MCS in the tethering group was 52.7 and for PCS was 46.8. The fusion group had MCS 52.3 and PCS 47.1. At the latest follow-up, the average SF-36 MCS in the tethering group was 56.9 and PCS 57.2. Fusion group had MCS 52.3 and PCS 53.1.

---

Federal and State law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. The member’s contract benefits in effect on the date that services are rendered must be used. Medical Policy, which addresses medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

**Scoliosis Surgery**

---

A retrospective chart review by Meyers in 2021 reports on the complications of 184 participants 90 days post-op vertebral body tethering. There were 12 participants who experienced 12 complications in this 90-day period following surgery. There were 6 participants with major complications including 3 chylothoraces, 2 hemothoraces, and 1 lumbar radiculopathy. Minor complications were experienced by 6 participants which included requirement of supplemental oxygen, superficial wound infection, prolonged nausea, and Raynaud phenomenon.

Broken tethers can lead to a loss of correction. However, some breakages may not have a negative effect on curve correction. A 2022 retrospective review by Trobisch and colleagues analyzed tether breakage rates following vertebral body tethering and the influence of breakage on curve behavior. There were 10 participants who had surgical revisions with 15 curves and 80 segments re-evaluated in vivo. Tether breakage was found in 36 segments. Of the 36 segments, 21 were suspected to have a tether breakage prior to revision surgery. During revision surgery 20 of 21 suspected breakages were confirmed. One suspected breakage measured an angular difference of 6.2° but was not found to be broken intraoperatively. Of the segments with an intraoperative confirmed breakage, 3 of them had a decreased segmental Cobb angle which was felt most likely due to growth modulation before the breakage occurred. For all confirmed cases of breakage, the average segmental loss of correction was 6.6°. While broken tethers are a potential complication of vertebral body tethering surgery, not all breakages have a negative impact on the correction.

In a 2021 retrospective review by Baroncini and colleagues, the authors measured pulmonary function before and after vertebral body tethering surgery of 51 participants at 6 weeks, 6 months, and 12 months postoperatively. Parameters measured were total lung capacity (TLC), forced expiratory volume in 1 second (FEV1), and forced vital capacity (FVC). TLC preoperatively was 98% ± 15, at 6 weeks was 93% ± 15, at 6 months was 96% ± 17, and at 12 months was 99% ± 15. FEV1 preoperatively was 85% ± 16, at 6 weeks was 80% ± 14, at 6 months was 84% ± 14, and at 12 months was 89% ± 9. FVC preoperatively was 91% ± 17, at 6 weeks was 90% ± 17, at 6 months was 90% ± 16, and at 12 weeks was 86% ± 9. The authors note that longer follow-up is necessary to evaluate the long-term effect of vertebral body tethering on pulmonary function and a comparison of pulmonary function between fusion surgeries and vertebral body tethering surgeries should be done to determine whether tethering is as beneficial as fusion in respect to pulmonary function.

In 2020, the Pediatric Orthopaedic Society of North American along with the Scoliosis Research Society released a position statement on payor coverage for vertebral body tethering for skeletally immature individuals with idiopathic scoliosis. The recommendation is “payors should provide coverage for any FDA approved devices under FDA stated clinical indications and requirements.” However, it should be noted these recommendations are not based on robust trials.

**Minimally Invasive Deformity Correction System**

A new proposed fusionless, surgical correction treatment of scoliosis is a minimally invasive deformity correction system.

In a 2021 retrospective review, Floman and colleagues sought to determine if a motion-sparing posterior device could modulate growth in skeletally immature individuals with adolescent idiopathic scoliosis. There were 45 individuals evaluated, who had a follow-up of at least 2 years following surgery. The mean pre-operative curve was 46°. There were 16 participants in the Risser 0-1 stage, 15 participants in the Risser 2-3 stage, and 14 participants in the Risser 4-5 stage at the time of surgery. There were 35 participants with Lenke type 1 curves and 10 participants with Lenke type 5 curves. The average preoperative major curve magnitude, of both curve types, was similar

---

Federal and State law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. The member's contract benefits in effect on the date that services are rendered must be used. Medical Policy, which addresses medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

**Scoliosis Surgery**

---

among the three Risser groups, 47.6°, 46° and 41.5°. At final follow-up, curves were reduced to 26.4°, 20.4° and 26.2°, respectively. Thoracic kyphosis increased by 7° on average in the Lenke 1 curves and lumbar lordosis decreased by 4° in the Lenke 5 curves. Revision surgery was required by 4 participants, 1 was converted to vertebral body tethering and in 3 cases there was a malfunction in pedicle screws, nut loosening, or ratchet malfunction. None of the revision surgeries were converted to a spinal fusion procedure.

A 2021 prospective, industry sponsored study by Stadhouder and colleagues reported on the use of the minimally invasive deformity correction system in 20 participants with adolescent idiopathic scoliosis. Inclusion criteria were 12 to 17 years of age, skeletal immaturity (defined as Risser stage 1 to 4), a single structural curve (Lenke type 1 or 5), a major Cobb angle of 40° to 55°, reduction of the major curve to less than 35° on x-ray, and apical vertebral rotation of less than 15°. With an average of 12 months of follow-up postoperatively, 7 of 20 participants showed signs of surgical failure and study recruitment was stopped. Subsequently, another 3 participants showed implant failures at further follow-up. Reasons for implant failures included pain, breakage/failure and osteolysis. Six of the participants required removal of the implant, 2 participants had revision to a new implant, and 2 participants had revision to posterior spinal fusion. The 10 remaining participants were followed for a mean of 3.8 years. The mean major curve for this group measured 45.4° preoperatively, 31.4° at 2 weeks postoperatively, and 31.0° at the time of the latest follow-up. The mean minor curve measured 31.3° preoperatively, 26.1° at 2 weeks postoperatively, and 24.2° at the time of the latest follow-up. There was no significant change in major or minor curve deformity after 2 weeks postoperatively. Lumbar lordosis and thoracic kyphosis remained unaltered and there were no changes in apical vertebral rotation. The authors concluded the rate of implant-related complications along with no correction of the curve or distraction of the ratchet observed postoperatively was rationale for early study termination.

Currently there is a paucity of literature depicting reasonable conclusions on health outcomes. There is also a lack of conclusion that this technique is as beneficial as established alternative treatments.

**Vertebral Body Stapling**

Another proposed alternative to bracing in the treatment of scoliosis is vertebral body stapling. The staples are surgically inserted into the vertebrae of the individual and designed to prevent further curvature of the spine.

Betz and colleagues (2003) reported the results of a study to determine the efficacy of vertebral body stapling in 21 individuals (27 curves) with adolescent idiopathic scoliosis. No major, but three instances of minor complications were noted. One individual experienced an intraoperative segmental vein bleed which resulted in an estimated blood loss of 1500 cc as compared to the average estimated blood loss of 247 cc for all participants. One subject developed a chylothorax and another developed pancreatitis. None of the individuals experienced staple dislodgement or movement during the follow-up period (mean 11 months, range 3-36 months), and no adverse effects specifically related to the staples were identified. Utility (defined as curve stability) was evaluated in 10 individuals with stapling with greater than 1-year follow-up (mean 22.6 months) and preoperative curve less than 50 degrees. Treatment failure was considered progression of greater than or equal to 6 degrees or beyond 50 degrees. Of these 10 individuals, 4 (40%) progressed and 6 (60%) remained stable or improved. One of 10 (10%) in the stapling group had progressed beyond 50 degrees and underwent spinal fusion. Six of the subjects required stapling of a second curve, 3 as part of the primary surgery, and 3 as a second stage because a second untreated curvature progressed. The authors concluded that vertebral body stapling for the treatment of scoliosis in adolescents was feasible and safe in this group of 21 subjects. However, the results need to be considered with caution, inasmuch as the follow-up period was short and there was no comparison of this technique with conventional treatment, such as bracing.

---

Federal and State law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. The member's contract benefits in effect on the date that services are rendered must be used. Medical Policy, which addresses medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

In 2005, Betz and colleagues carried forward their clinical series and presented the retrospective findings of 39 consecutive individuals (52 curves) who received vertebral body stapling as treatment for idiopathic scoliosis or scoliosis associated with other conditions, such as Marfan syndrome or skeletal dysplasia (syndromic scoliosis). Complications were reported in 6 cases. A 4-year-old with infantile idiopathic scoliosis developed a rupture of a pre-existing undiagnosed diaphragmatic hernia which required emergency repair. One participant experienced a puncture in a segmental spinal vein secondary to a staple prong and required both transfusion and conversion to an open procedure to control blood loss. One subject developed chylothorax as a result of a staple puncture of the thoracic duct at T12. This individual was treated with a chest tube and total parenteral nutrition. Another participant experienced mild pancreatitis. Clinically significant atelectasis was experienced by 2 individuals and 2 other participants required prolonged chest tube drainage (greater than 4 days). In 31 subjects who were followed for an average of 12 months, there were no reports of staple dislodgement or migration. However, there was one report of staple fracture. Five participants (15%) progressed during follow-up and required spinal fusion.

In another study, Betz and colleagues (2010) reported the findings of vertebral body stapling in 28 individuals with idiopathic scoliosis for a minimum follow-up period of at least 2 years. The authors reported a success rate (curves corrected to within 10 degrees of preoperative measurement or decreased > 10 degrees) in 87% of all of the lumbar curves and 77% in thoracic curves measuring less than 35 degrees. In the cases of thoracic curves, which measured greater than 35 degrees, vertebral body stapling was not considered successful and required alternative treatments. In the conclusions section of the article, the authors acknowledged the limitations of the study and cautioned the reader that the "results should be considered preliminary as follow-up to skeletal maturity will be needed before definitive results can be described."

Laituri and colleagues (2013) reported the results of a retrospective study on children who underwent thoracoscopic vertebral body stapling for juvenile scoliosis from January 2007 to December 2010. Only individuals with a follow-up of at least 2 years were included in this study group. Data considered were demographics, indications for vertebral body stapling, degree of curvature, treatment, complications, and follow-up. Cobb angle was used to measure the initial degree of curvature on a standing posterior-anterior spine X-Ray. During the study period, 11 individuals underwent thoracoscopic vertebral body stapling for juvenile idiopathic scoliosis using single lung ventilation in a lateral position. The study group consisted of 7 subjects between the ages of 8-11 years with at least a 2-year follow-up. Indications for stapling in these 7 participants were progression of scoliosis (n=3), noncompliance with brace (n=3), and double curve with progression (n=1). The mean preoperative Cobb angle was  $34.1 \pm 5^\circ$  (range, 25-41°), and the mean immediate postoperative Cobb angle measurement was  $23 \pm 5^\circ$  (range, 16-30°). The staples encompassed a mean number of 6.4 vertebral bodies. The mean duration of chest drainage was 2.7 days, the mean length of hospitalization was 3.9 days and the mean operative time was  $156.2 \pm 39.5$  minutes. The authors indicated there were no intraoperative complications or mortality. During the postoperative period, 1 individual developed a pleural effusion on the contralateral side that required drainage. These 7 participants were followed for a mean of 34 months (range, 29-44 months). The mean Cobb angle at last follow-up was  $24.7^\circ$  (range, 15-38°). At the time of last follow-up, none of the participants required postoperative bracing or spinal fusion. The authors concluded that thoracoscopic vertebral body stapling is a safe and effective method for treating progressive scoliosis in young children.

Theologis and colleagues (2013) evaluated 12 females older than 10 years of age with idiopathic thoracic or lumbar scoliosis of 30° to 39° who were treated with vertebral body stapling. The participants were followed for a minimum of 24 months. Outcome variables included curve progression and magnitude, surgical complications and

---

Federal and State law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. The member's contract benefits in effect on the date that services are rendered must be used. Medical Policy, which addresses medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

a need for reoperation. The preoperative and postoperative curve magnitudes were compared. A total of 13 curves were treated with vertebral body stapling (lumbar: n=4, thoracic: n=9). The follow-up period ranged from 2.2-5.4 years and averaged 3.4 years. The average preoperative curve magnitude was 33.4° (range, 30-39°) compared to most recent curve magnitude measurement at follow-up of 23.0° (range, 10-34°). All curves, both thoracic and lumbar, were treated successfully. Postoperative curve magnitudes did not change significantly between the first erect radiographs and the most recent follow-up. Two of the study participants had pneumothorax, and 1 participant had symptomatic pleural effusion. None of the study participants required definitive fusion for curve progression. The authors concluded that vertebral body stapling is an effective method to control curve progression in the high-risk group of children younger than 10 years with idiopathic scoliosis between 30° and 39° in whom bracing may be ineffective.

In 2018, Cahill and colleagues performed a retrospective review on 63 subjects between 7 to 15 years of age with idiopathic scoliosis. The aim of this study was to evaluate the change in Cobb angle measurements over time in subjects treated with vertebral body stapling. Outcomes were assessed by using three categories. Cahill (2018) stated:

“Improvement” was defined as a decrease in the preoperative Cobb angle of greater than 10°. “No change” was defined as a +10° to -10° change in the preoperative Cobb angle (both values inclusive). “Progression” was defined as an increase of the curve by greater than 10°. These assessments allowed for the classification of success versus failure, with “success” defined as either improvement or no change and “failure” defined as progression.

The authors reported that of the subjects who had vertebral body stapling of the lumbar curve, 82% were successful, and of the subjects who had vertebral body stapling of the thoracic curve, 74% were successful.

A retrospective chart review by Trupia and colleagues (2019) reported on 10 skeletally immature participants with adolescent idiopathic scoliosis who underwent vertebral body stapling. The participants had curves ranging from 25° to 35° prior to surgery. The average duration of follow-up was 6.4 years. At the first postoperative visit, all participants showed curve correction. At the final follow-up visit, half of the participants showed curve progression greater than 5° while the other half of the participants either remained stable or corrected over time. The 5 participants who showed curve progression were younger than those who remained stable (10.8 years versus 12.8 years respectively). Four of these participants required further surgery for worsening of scoliosis. Three participants had hardware-related complications including breaking of a distal staple and asymptomatic loosening of a staple. None of the complications required further intervention. Limitations include the retrospective design. The authors state “In light of these results and the potential for surgical and hardware-related complications, we no longer recommend vertebral stapling, regardless of curve size or skeletal maturity.”

In a 2020 retrospective case series, Murray and colleagues reported on 7 children with juvenile idiopathic scoliosis who underwent vertebral body stapling. Using radiologic imaging, the aim of the study was to measure the rate of growth of vertebral bodies for 6 years following surgery. The average preoperative Cobb angle was 30°, with a decrease to 20° at the first postoperative visit. One participant has shown an improvement of greater than 10°, 4 participants have shown no change in their curve, and 2 participants have shown progression of their curves by more than 10°. Average growth rate for all participants was 0.86 mm/year per vertebral body on the side which was stapled compared to 0.83 mm/year per vertebral body on the unstapled side. There was no significant difference in

---

Federal and State law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. The member’s contract benefits in effect on the date that services are rendered must be used. Medical Policy, which addresses medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.



**Scoliosis Surgery**

the growth rate of the vertebral bodies between the stapled and unstapled sides and the authors conclude “the staple does not generate sufficient force to modulate growth.”

**Summary**

While the use of a motion-sparing posterior device may show promise for treatment of adolescent idiopathic scoliosis, at this time there is a lack of published peer-reviewed medical literature which shows reasonable conclusions on health outcomes. The clinical evidence on vertebral body stapling is not robust enough to make determinations regarding its safety and efficacy. There is still considerable risk of curve progression for these study participants and it may be premature to conclude that vertebral body stapling is an effective means of controlling curve progression in high-risk individuals who have not reached skeletal maturity. Study results once the individuals have reached skeletal maturity are warranted in order to determine the definitive benefit of vertebral body stapling in these individuals at high risk for continued curve progression.

**Background/Overview****Vertebral Body Tethering**

Vertebral body tethering is a technique in which bone screws are anchored to the front of each vertebral bone in the curved area of the spinal column. A flexible cord, or tether, is attached to the screws and tensioned to attain the desired degree of spine straightening. In August 2019, the FDA granted HDE for The Tether™ Vertebral Body Tethering System (Zimmer Biomet Spine, Inc., Westminster, CO). The device is indicated for skeletally immature individuals with idiopathic scoliosis (major Cobb angle of 30 to 65 degrees) who have failed bracing and/or are intolerant to brace wear. Some of the benefits to vertebral body tethering include allowing for continued growth and mobility, faster recovery time, spinal motion sparing, and less placement of hardware. However, concerns for vertebral body tethering include the possibility of overcorrection of the curve, potential disc degeneration within the instrumented spine, potential for fixation failure or cord breakage, and infection.

**Minimally Invasive Deformity Correction System**

The minimally invasive deformity correction system is a fusion-less surgical treatment for scoliosis. It is a ratchet-based, expandable rod which attaches to the spine using pedicle screws. This internal brace, like the vertebral body tethering and stapling, achieves correction without the need for spinal fusion. It is implanted posteriorly on one side of the spine. An example is the ApiFix system which received HDE from the FDA in 2019. The device is indicated for treatment of individuals with adolescent idiopathic scoliosis. The Summary of Safety and Probable Benefit document by the FDA lists potential adverse events associated with the use of the device some of which include screw/nut loosening, device migration or breakage, and inadequate curve correction or loss of curve correction.

**Vertebral Body Stapling**

Vertebral body stapling is being studied as an alternative to bracing or spinal fusion for the treatment of progressive idiopathic scoliosis in skeletally immature individuals. Because this procedure avoids fusion of the spine, it is proposed that this treatment will permit a gradual correction of the spinal curvature as the individual grows while maintaining movement and flexibility and decreasing the risk for back pain in adulthood. Benefits include it is believed to be more comfortable and less embarrassing than wearing a brace. Unlike spinal fusion, stapling offers the advantage of allowing the individual to retain the flexibility of their spine. However, complications involving the staples may include breakage, loosening, or dislodging.

**Definitions**

Federal and State law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. The member's contract benefits in effect on the date that services are rendered must be used. Medical Policy, which addresses medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

**Scoliosis Surgery**

**Cobb angle:** Determined by radiographs, a standard measurement to diagnose and track progression of scoliosis. The angle is determined by the intersection of tangential lines which are drawn from the superior end plate of the superior vertebra and the inferior end plate of the inferior vertebra.

**Curve flexibility:** The percentage of change in Cobb angle from standing upright images to flexibility images.

**Risser score:** A grading system used to measure ossification of the iliac apophysis.

**Sanders hand score:** A predictor of the curve acceleration phase of growth based on an x-ray of the left hand.

**Scoliosis:** A spinal disorder characterized by abnormal lateral curvature of the spine and vertebral rotation.

**Coding**

*The following codes for treatments and procedures applicable to this document are included below for informational purposes. Inclusion or exclusion of a procedure, diagnosis or device code(s) does not constitute or imply member coverage or provider reimbursement policy. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.*

*Vertebral body tethering*

**When services may be Medically Necessary when criteria are met:**

**CPT**

- 0656T Vertebral body tethering, anterior; up to 7 vertebral segments
- 0657T Vertebral body tethering, anterior; 8 or more vertebral segments

**ICD-10 Procedure**

- 0PS403Z Reposition thoracic vertebra with spinal stabilization device, vertebral body tether, open approach
- 0PS443Z Reposition thoracic vertebra with spinal stabilization device, vertebral body tether, percutaneous endoscopic approach
- 0QS003Z Reposition lumbar vertebra with spinal stabilization device, vertebral body tether, open approach
- 0QS043Z Reposition lumbar vertebra with spinal stabilization device, vertebral body tether, percutaneous endoscopic approach

**ICD-10 Diagnosis**

- M41.00-M41.9 Scoliosis
- Q67.5 Congenital deformity of spine (congenital scoliosis NOS)

**When services are Investigational and Not Medically Necessary:**

For the procedure codes listed above when criteria are not met, or for all other diagnoses not listed.

*Other procedures*

**When services are Investigational and Not Medically Necessary:**

Federal and State law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. The member's contract benefits in effect on the date that services are rendered must be used. Medical Policy, which addresses medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

**Scoliosis Surgery**

When the code describes a procedure indicated in the Position Statement section as investigational and not medically necessary.

**CPT**

22899 Unlisted procedure, spine [when specified as vertebral body stapling or implantation of a posterior (dynamic) distraction device]

**ICD-10 Procedure**

For the following codes, when specified as vertebral body stapling:  
 0PH404Z Insertion of internal fixation device into thoracic vertebra, open approach  
 0PH434Z Insertion of internal fixation device into thoracic vertebra, percutaneous approach  
 0PH444Z Insertion of internal fixation device into thoracic vertebra, percutaneous endoscopic approach  
 0QH004Z Insertion of internal fixation device into lumbar vertebra, open approach  
 0QH034Z Insertion of internal fixation device into lumbar vertebra, percutaneous approach  
 0QH044Z Insertion of internal fixation device into lumbar vertebra, percutaneous endoscopic approach

**ICD-10 Diagnosis**

M41.00-M41.9 Scoliosis  
 Q67.5 Congenital deformity of spine (congenital scoliosis NOS)

**When services are also Investigational and Not Medically Necessary:**

**ICD-10 Procedure**

XNS00C7 Reposition of lumbar vertebra using posterior (dynamic) distraction device, open approach, new technology group 7  
 XNS03C7 Reposition of lumbar vertebra using posterior (dynamic) distraction device, percutaneous approach, new technology group 7  
 XNS40C7 Reposition of thoracic vertebra using posterior (dynamic) distraction device, open approach, new technology group 7  
 XNS43C7 Reposition of thoracic vertebra using posterior (dynamic) distraction device, percutaneous approach, new technology group 7

**ICD-10 Diagnosis**

All diagnoses

**References**

**Peer Reviewed Publications:**

1. Baker CE, Kiebzak GM, Neal KM. Anterior vertebral body tethering shows mixed results at 2-year follow-up. *Spine Deform.* 2021; 9(2):481-489.
2. Betz RR, D'Andrea LP, Mulcahey MJ, Chafetz RS. Vertebral body stapling procedure for the treatment of scoliosis in the growing child. *Clin Orthop Relat Res.* 2005; (434):55-60.

Federal and State law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. The member's contract benefits in effect on the date that services are rendered must be used. Medical Policy, which addresses medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

**Scoliosis Surgery**

3. Betz RR, Kim J, D'Andrea LP, et al. An innovative technique of vertebral body stapling for the treatment of patients with adolescent idiopathic scoliosis: a feasibility, safety, and utility study. *Spine (Phila Pa 1976)*. 2003; 28(20):S255-S265.
4. Betz RR, Ranade A, Samdani AF, et al. Vertebral body stapling: a fusionless treatment option for a growing child with moderate idiopathic scoliosis. *Spine (Phila Pa 1976)*. 2010; 35(2):169-176.
5. Baroncini A, Trobisch P, Blau C, et al. Analysis of the pulmonary function in patients undergoing vertebral body tethering for adolescent idiopathic scoliosis. *Eur Spine J*. 2021 Oct 22. Online ahead of print.
6. Cahill PJ, Auriemma M, Dakwar E, et al. Factors predictive of outcomes in vertebral body stapling for idiopathic scoliosis. *Spine Deform*. 2018; 6(1):28-37.
7. Floman Y, El-Hawary R, Lonner BS, et al. Vertebral growth modulation by posterior dynamic deformity correction device in skeletally immature patients with moderate adolescent idiopathic scoliosis. *Spine Deform*. 2021; 9(1):149-153.
8. Guille JT, D'Andrea LP, Betz RR. Fusionless treatment of scoliosis. *Orthop Clin North Am*. 2007; 38(4):541-545, vii.
9. Hoernschemeyer DG, Boeyer ME, Robertson ME, et al. Anterior vertebral body tethering for adolescent scoliosis with growth remaining: a retrospective review of 2 to 5-year postoperative results. *J Bone Joint Surg Am*. 2020; 102(13):1169-1176.
10. Laituri CA, Schwend RM, Holcomb GW 3rd. Thoracoscopic vertebral body stapling for treatment of scoliosis in young children. *J Laparoendosc Adv Surg Tech A*. 2012; 22(8):830-833.
11. Meyers J, Eaker L, von Treuheim TDP, et al. Early operative morbidity in 184 cases of anterior vertebral body tethering. *Sci Rep*. 2021; 11(1):23049.
12. Miyajiri F, Pawelek J, Nasto LA, et al. Safety and efficacy of anterior vertebral body tethering in the treatment of idiopathic scoliosis. *Bone Joint J*. 2020; 102-B(12):1703-1708.
13. Murray E, Tung R, Sherman A, Schwend RM. Continued vertebral body growth in patients with juvenile idiopathic scoliosis following vertebral body stapling. *Spine Deform*. 2020; 8(2):221-226.
14. Newton PO, Bartley CE, Bastrom TP, et al. Anterior spinal growth modulation in skeletally immature patients with idiopathic scoliosis: a comparison with posterior spinal fusion at 2 to 5 years postoperatively. *J Bone Joint Surg Am*. 2020; 102(9):769-777.
15. Newton PO, Kluck DG, Saito W, et al. Anterior spinal growth tethering for skeletally immature patients with scoliosis: a retrospective look two to four years postoperatively. *J Bone Joint Surg Am*. 2018; 100(19):1691-1697.
16. Pehlivanoglu T, Oltulu I, Erdag Y, et al. Comparison of clinical and functional outcomes of vertebral body tethering to posterior spinal fusion in patients with adolescent idiopathic scoliosis and evaluation of quality of life: preliminary results. *Spine Deform*. 2021; 9(4):1175-1182.
17. Pehlivanoglu T, Oltulu I, Ofluoglu E, et al. Thoracoscopic vertebral body tethering for adolescent idiopathic scoliosis: a minimum of 2 years' results of 21 patients. *J Pediatr Orthop*. 2020; 40(10):575-580.
18. Rushton PRP, Nasto L, Parent S, et al. Anterior vertebral body tethering (AVBT) for treatment of idiopathic scoliosis in the skeletally immature: results of 112 cases. *Spine (Phila Pa 1976)*. 2021; 46(21):1461-1467.
19. Samdani AF, Ames RJ, Kimball JS, et al. Anterior vertebral body tethering for idiopathic scoliosis: two-year results. *Spine (Phila Pa 1976)*. 2014; 39(20):1688-1693.
20. Samdani AF, Ames RJ, Kimball JS, et al. Anterior vertebral body tethering for immature adolescent idiopathic scoliosis: One-year results on the first 32 patients. *Eur Spine J*. 2015; 24(7):1533-1539.
21. Samdani AF, Pahys JM, Ames RJ, et al. Prospective follow-up report on anterior vertebral body tethering for idiopathic scoliosis: interim results from an FDA IDE study. *J Bone Joint Surg Am*. 2021; 103(17):1611-1619.

Federal and State law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. The member's contract benefits in effect on the date that services are rendered must be used. Medical Policy, which addresses medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

**Scoliosis Surgery**

22. Shin M, Arguelles GR, Cahill PJ, et al. Complications, reoperations, and mid-term outcomes following anterior vertebral body tethering versus posterior spinal fusion: A Meta-Analysis. *JB JS Open Access*. 2021; 6(2):e21.00002.
23. Stadhouders A, Holewijn RM, Haanstra TM, et al. High failure rates of a unilateral posterior peri-apical distraction device (ApiFix) for fusionless treatment of adolescent idiopathic scoliosis. *J Bone Joint Surg Am*. 2021; 103(19):1834-1843.
24. Theologis AA, Cahill P, Auriemma M, et al. Vertebral body stapling in children younger than 10 years with idiopathic scoliosis with curve magnitude of 30° to 39°. *Spine*. 2013; 38(25):E1583-E1588.
25. Torre-Healy A, Samdani AF. Newer technologies for the treatment of scoliosis in the growing spine. *Neurosurg Clin N Am*. 2007; 18(4):697-705.
26. Trobisch P, Baroncini A, Berrer A, Da Paz S. Difference between radiographically suspected and intraoperatively confirmed tether breakages after vertebral body tethering for idiopathic scoliosis. *Eur Spine J*. 2022 Jan 8. Online ahead of print.
27. Trupia E, Hsu AC, Mueller JD, et al. Treatment of idiopathic scoliosis with vertebral body stapling. *Spine Deform*. 2019; 7(5):720-728.
28. Wong HK, Ruiz JNM, Newton PO, Gabriel Liu KP. Non-fusion surgical correction of thoracic idiopathic scoliosis using a novel, braided vertebral body tethering device: minimum follow-up of 4 years. *JB JS Open Access*. 2019; 4(4):e0026.

**Government Agency, Medical Society, and Other Authoritative Publications:**

1. Pediatric Orthopaedic Society of North America. Joint SRS/POSNA position statement on payor coverage for anterior fusionless scoliosis technologies for immature patients with idiopathic scoliosis. 2020. Available at: <https://posna.org/getattachment/Physician-Education/Position-Statements/Why-Should-Insurance-Cover-AVBT-April-2020.pdf?lang=en-US>. Accessed on February 25, 2022.
2. U.S. Food and Drug Administration (FDA) 510 (k) Premarket Notification Summary. OSStaple™ Staple System. No. K061385. Rockville, MD. FDA. May 30, 2007. Available at: [http://www.accessdata.fda.gov/cdrh\\_docs/pdf6/K061385.pdf](http://www.accessdata.fda.gov/cdrh_docs/pdf6/K061385.pdf). Accessed on February 25, 2022.
3. U.S. Food and Drug Administration (FDA) HDE Notification Summary. The Tether™ Vertebral Body Tethering System. No. H190005. Rockville, MD. FDA. August 16, 2019. Available at: [https://www.accessdata.fda.gov/cdrh\\_docs/pdf19/H190005A.pdf](https://www.accessdata.fda.gov/cdrh_docs/pdf19/H190005A.pdf). Accessed on February 25, 2022.
4. U.S. Food and Drug Administration (FDA) Summary of Safety and Probable Benefit. Minimally Invasive Deformity Correction (MID-C) System. No. H170001. Rockville, MD. FDA. August 23, 2019. Available at: [https://www.accessdata.fda.gov/cdrh\\_docs/pdf17/H170001B.pdf](https://www.accessdata.fda.gov/cdrh_docs/pdf17/H170001B.pdf). Accessed on February 25, 2022.
5. Zimmer Biomet. Safety and efficacy study of spinal tethering. NLM Identifier: NCT02897453. Last updated on August 20, 2020. Available at: <https://www.clinicaltrials.gov/ct2/show/NCT02897453?cond=NCT02897453&rank=1>. Accessed on February 25, 2022.

**Websites for Additional Information**

1. American Academy of Family Physicians. Adolescent idiopathic scoliosis: diagnosis and management. Available at: <https://www.aafp.org/afp/2014/0201/afp20140201p193.pdf>. Accessed on March 24, 2022.

**Index**

**ApiFix**

Federal and State law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. The member’s contract benefits in effect on the date that services are rendered must be used. Medical Policy, which addresses medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

**Scoliosis Surgery**

Minimally Invasive Deformity Correction System  
 Nitinol Staple  
 OSStaple™  
 The Tether™ Vertebral Body Tethering System

**The use of specific product names is illustrative only. It is not intended to be a recommendation of one product over another, and is not intended to represent a complete listing of all products available.**

**Document History**

Status	Date	Action
Revised	05/12/2022	Medical Policy & Technology Assessment Committee (MPTAC) review. Added MN criteria for vertebral body tethering. Added Definitions and Websites for Additional Information sections. Updated Rationale, Coding and References sections.
Revised	11/11/2021	MPTAC review. Title changed to Scoliosis Surgery. Added minimally invasive deformity correction system to the scope of the document and Position Statement. Updated Description/Scope, Rationale, Background/Overview, References, and Index sections. Updated Coding section; added codes XNS00C7, XNS03C7, XNS40C7, XNS43C7.
	07/01/2021	Updated Coding section with 07/01/2021 CPT changes and 10/01/2021 ICD-10-PCS changes; added 0656T, 0657T and 0PS403Z, 0PS443Z, 0QS003Z, 0QS043Z.
Reviewed	11/05/2020	MPTAC review. Updated Rationale, References, and Index sections.
Revised	11/07/2019	MPTAC review. Revised scope of document to include vertebral body tethering. Title changed. Added vertebral body tethering to INV/NMN statement. Updated Description/Scope, Rationale, Background/Overview, References, and Index sections.
Reviewed	01/24/2019	MPTAC review.
Reviewed	01/25/2018	MPTAC review. The document header wording updated from “Current Effective Date” to “Publish Date.” Updated Rationale and References sections.
Reviewed	02/02/2017	MPTAC review. Updated the review date, References and History sections of the document.
Reviewed	02/04/2016	MPTAC review. Updated the review date, References and History sections of the document. Removed ICD-9 codes from Coding section.
Reviewed	02/05/2015	MPTAC review. Updated the review date, Description/Scope, References and History sections of the document.
Reviewed	02/13/2014	MPTAC review. Updated the review date, Rationale, References and History sections of the document.
Reviewed	02/14/2013	MPTAC review. Updated the review date, References and History sections of the document.
Reviewed	02/16/2012	MPTAC review. Updated the review date, References and History sections of the document.
Reviewed	02/17/2011	MPTAC review. Updated the review date, Rationale, References and History sections of the document.

Federal and State law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. The member’s contract benefits in effect on the date that services are rendered must be used. Medical Policy, which addresses medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

**Scoliosis Surgery**

---

Reviewed	02/25/2010	MPTAC review. Updated the review date, Description/Scope, Rationale, Background/Overview, References and History sections of the document.
Reviewed	02/26/2009	MPTAC review. Changed title to “Vertebral Body Stapling for the Treatment of Scoliosis in Children and Adolescents.” Revised Position Statement to indicate that vertebral body stapling is investigational and not medically necessary as a treatment of scoliosis in both children and adolescents. Updated review date, Rationale, Background/Overview, References and History sections of the document.
Reviewed	02/21/2008	MPTAC review. Updated review date, References and History sections of the document. The phrase “investigational/not medically necessary” was clarified to read “investigational and not medically necessary.” This change was approved at the November 29, 2007 MPTAC meeting.
New	03/08/2007	MPTAC review. Initial document development.

---

Federal and State law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. The member’s contract benefits in effect on the date that services are rendered must be used. Medical Policy, which addresses medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.